antibodies - online.com







anti-SLIT1 antibody (AA 308-512)



Images



Overview

Quantity:	100 μL
Target:	SLIT1
Binding Specificity:	AA 308-512
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLIT1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Purpose:	Polyclonal Antibody to Slit Homolog 1 (Slit1)
Immunogen:	Recombinant Slit Homolog 1 (Slit1) corresdonding to Pro308~Ala512 (Accession # 088279)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against Slit1. It has been selected for its ability to recognize Slit1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Expiry Date:


24 months

	Target:
Products)	Alternative Name:
Multiple epidermal growth factor-like domains protein 4	Background:
	Pathways:
	Application Details
2 μg/mL Immunocytochemistry in formalin fixed cells: 5-20 μg/mL ry in formalin fixed frozen section: 5-20 μg/mL Immunohistochemis μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optimast be determined by end user.	Application Notes:
s described by the loss rate. The loss rate was determined by accelerate, that is, incubate the protein at 37°C for 48h, and no obvious ipitation were observed. The loss rate is less than 5% within the expire storage condition.	Comment:
у	Restrictions:
	Handling
	Format:
ontaining 0.05 % Proclin-300, 50 % glycerol.	Buffer:
	Preservative:
ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which shou	Precaution of Use:
	Storage:
ent use. Stored at -20°C in a manual defrost freezer for two year with ivity. Avoid repeated freeze-thaw cycles.	Storage Comment:
s described by the loss rate. The loss rate was determined by accesest, that is, incubate the protein at 37°C for 48h, and no obvious ipitation were observed. The loss rate is less than 5% within the experimental strange condition. The loss rate is less than 5% within the experimental strange condition. The loss rate is less than 5% within the experimental strange condition. The loss rate is less than 5% within the experimental strange condition. The loss rate is less than 5% within the experimental strange condition. The loss rate is less than 5% within the experimental strange condition. The loss rate is less than 5% within the experimental strange condition. The loss rate is less than 5% within the experimental strange condition.	Restrictions: Handling Format: Buffer: Preservative: Precaution of Use: Storage:




Immunohistochemistry

Image 1. Detection of Slit1 in Rat Adrenal gland Tissue using Polyclonal Antibody to Slit Homolog 1 (Slit1)



Western Blotting

Image 2. Detection of Recombinant Slit1, Rat using Polyclonal Antibody to Slit Homolog 1 (Slit1)



Immunohistochemistry

Image 3. Detection of Slit1 in Rat Heart Tissue using Polyclonal Antibody to Slit Homolog 1 (Slit1)

Please check the product details page for more images. Overall 7 images are available for ABIN7438097.