antibodies - online.com







anti-Periostin antibody (AA 370-494)

Images



Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | Periostin (POSTN) |
| Binding Specificity: | AA 370-494 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Periostin antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| Purpose: | Polyclonal Antibody to Periostin (POSTN) |
|-------------------|--|
| Immunogen: | Recombinant Periostin (POSTN) corresdonding to Asp370~lle494 with N-terminal His Tag |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against POSTN. It has been selected for its ability to recognize POSTN in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Human, Rat |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |


Target Details


Expiry Date:

| Target Details | |
|---------------------|---|
| Target: | Periostin (POSTN) |
| Alternative Name: | Periostin (POSTN Products) |
| Background: | OSF-2, PDLPOSTN, PN, Osteoblast Specific Factor 2 |
| Application Details | |
| Application Notes: | Western blotting: 0.5-2 μ g/mL Immunohistochemistry: 5-20 μ g/mL Immunocytochemistry: 5-20 μ g/mL Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without |

detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months





Western Blot 44 33 26 22 18 -15kDa 14 10

Western Blotting

Image 1. Detection of POSTN in BXPC3 cell lysate using Polyclonal Antibody to Periostin (POSTN)

Western Blotting

Image 2. Detection of Recombinant POSTN, Mouse using Polyclonal Antibody to Periostin (POSTN)