# antibodies -online.com





## anti-CYP2E1 antibody (AA 34-273)



## **Images**



#### Go to Product pag

_					
	W	0	rv	10	W


Quantity:	100 μL	
Target:	CYP2E1	
Binding Specificity:	AA 34-273	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CYP2E1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

#### **Product Details**

Purpose:	Polyclonal Antibody to Cytochrome P450 2E1 (CYP2E1)	
Immunogen:	Recombinant Cytochrome P450 2E1 (CYP2E1) corresdonding to Pro34~Met273 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against CYP2E1. It has been selected for its ability to recognize CYP2E1 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

## **Target Details**

Application Details  Application Notes: Western blotting: 0.5-2 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemiparaffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optim working dilutions must be determined by end user.  Comment: The thermal stability is described by the loss rate. The loss rate was determined by acceethermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expediate under appropriate storage condition.  Restrictions: For Research Use only  Handling  Format: Liquid  Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative: ProClin  Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage: 4 °C,-20 °C			
Background:  CPE1, CYPIIE1, P450-J, P450C2E, 4-nitrophenol 2-hydroxylase, Cytochrome P450 Family Subfamily E1, Cytochrome P450, Subfamily IIE(Ethanol-Inducible) Polypeptide 1  Application Details  Application Notes:  Western blotting: 0.5-2 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in formalin fixed Immunosorbent Assay: 0.05-2 µg/mL Optim working dilutions must be determined by end user.  Comment:  The thermal stability is described by the loss rate. The loss rate was determined by acce thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expediate under appropriate storage condition.  Restrictions:  For Research Use only  Handling  Format:  Liquid  Buffer:  0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative:  ProClin  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage:  4 °C, 20 °C  Storage Comment:  Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year will detectable loss of activity. Avoid repeated freeze-thaw cycles.	Target:	CYP2E1	
Application Details  Application Notes: Western blotting: 0.5-2 μg/mL Immunocytochemistry in formalin fixed cells: 5-20 μg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 μg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 μg/mL Immunohistochemi paraffin section: 5-20 μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optim working dilutions must be determined by end user.  Comment: The thermal stability is described by the loss rate. The loss rate was determined by accee thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expediate under appropriate storage condition.  Restrictions: For Research Use only  Handling  Format: Liquid  Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative: ProClin  Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage: 4 °C,-20 °C  Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year will detectable loss of activity. Avoid repeated freeze-thaw cycles.	Alternative Name:	Cytochrome P450 2E1 (CYP2E1 Products)	
Application Notes:  Western blottling: 0.5-2 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optim working dilutions must be determined by end user.  Comment:  The thermal stability is described by the loss rate. The loss rate was determined by acce thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expediate under appropriate storage condition.  Restrictions:  For Research Use only  Handling  Format:  Liquid  Buffer:  0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative:  ProClin  Precaution of Use:  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage:  4 °C,-20 °C  Storage Comment:  Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Background:	CPE1, CYPIIE1, P450-J, P450C2E, 4-nitrophenol 2-hydroxylase, Cytochrome P450 Family 2 Subfamily E 1, Cytochrome P450,Subfamily IIE(Ethanol-Inducible)Polypeptide 1	
Immunohistochemistry in formalin fixed frozen section: 5-20 μg/mL Immunohistochemi paraffin section: 5-20 μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optim working dilutions must be determined by end user.  Comment: The thermal stability is described by the loss rate. The loss rate was determined by acce thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expediate under appropriate storage condition.  Restrictions: For Research Use only  Handling  Format: Liquid  Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative: ProClin  Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage: 4 °C,-20 °C  Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year wit detectable loss of activity. Avoid repeated freeze-thaw cycles.	Application Details		
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expediate under appropriate storage condition.  Restrictions:  For Research Use only  Handling  Format:  Liquid  Buffer:  0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative:  ProClin  Precaution of Use:  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage:  4 °C,-20 °C  Storage Comment:  Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Application Notes:	Immunohistochemistry in formalin fixed frozen section: 5-20 μg/mL Immunohistochemistry in paraffin section: 5-20 μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optimal	
Handling  Format: Liquid  Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative: ProClin  Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage: 4 °C,-20 °C  Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Comment:	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
Format:  Liquid  Buffer:  0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative:  ProClin  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage:  4 °C,-20 °C  Storage Comment:  Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Restrictions:	For Research Use only	
Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.  Preservative: ProClin  Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage: 4 °C,-20 °C  Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Handling		
Preservative:  ProClin  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage:  4 °C,-20 °C  Storage Comment:  Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Format:	Liquid	
Precaution of Use:  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sho handled by trained staff only.  Storage:  4 °C,-20 °C  Storage Comment:  Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
handled by trained staff only.  Storage: 4 °C,-20 °C  Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Preservative:	ProClin	
Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year with detectable loss of activity. Avoid repeated freeze-thaw cycles.	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
detectable loss of activity. Avoid repeated freeze-thaw cycles.	Storage:	4 °C,-20 °C	
Expiry Date: 24 months	Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
	Expiry Date:	24 months	




#### **Western Blotting**

**Image 1.** Detection of CYP2E1 in Human Hela cell lysate using Polyclonal Antibody to Cytochrome P450 2E1 (CYP2E1)

#### **Western Blotting**

Image 2. Detection of Recombinant CYP2E1, Mouse using Polyclonal Antibody to Cytochrome P450 2E1 (CYP2E1)



### **Immunohistochemistry**

Image 3. Detection of CYP2E1 in Mouse Liver Tissue using Polyclonal Antibody to Cytochrome P450 2E1 (CYP2E1)

Please check the product details page for more images. Overall 6 images are available for ABIN7438247.