antibodies - online.com







anti-HGFA antibody (AA 104-351)



Image



\sim			
	N/6	1//r	$I \cap V$

Quantity:	100 μL	
Target:	HGFA	
Binding Specificity:	AA 104-351	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HGFA antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details


Purpose:	Polyclonal Antibody to Hepatocyte Growth Factor Activator (HGFAC)	
Immunogen:	Recombinant Hepatocyte Growth Factor Activator (HGFAC) corresdonding to Pro104~Asp351 (Accession # Q9R098) with N-terminal His and GST Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against HGFAC. It has been selected for its ability to recognize HGFAC in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details

Expiry Date:

24 months

Target Details		
Target:	HGFA	
Alternative Name:	Hepatocyte Growth Factor Activator (HGFA Products)	
Background:	HGFA, HGFAP, HGF Activator	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	



Western Blotting

Image 1. Detection of Recombinant HGFAC, Mouse using Polyclonal Antibody to Hepatocyte Growth Factor Activator (HGFAC)