antibodies - online.com







anti-LMYC antibody (AA 186-439)





()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL
Target:	LMYC (MYCL)
Binding Specificity:	AA 186-439
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LMYC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Purpose:	Polyclonal Antibody to V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC)
Immunogen:	Recombinant V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC) corresdonding to Ala186~Ala439 (Accession # P01108) with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MYC. It has been selected for its ability to recognize MYC in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	LMYC (MYCL)
Alternative Name:	V-Myc Myelocytomatosis Viral Oncogene Homolog (MYCL Products)
Background:	C-Myc, BHLHE39, Myc Proto-Oncogene Protein, VMyc Class E basic helix-loop-helix protein 39, Transcription factor p64, Proto-oncogene c-Myc
Pathways:	Warburg Effect

Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	1:300-1200 Immunohistochemistry: 5-20 μg/mL
	1:30-120 Immunocytochemistry: 5-20 μg/mL
	1:30-120 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Western Blotting

Image 1. Detection of MYC in Raji cell lysate using Polyclonal Antibody to V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC)

Western Blotting

Image 2. Detection of MYC in HL60 cell lysate using Polyclonal Antibody to V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC)