# antibodies -online.com





# anti-N-Cadherin antibody (AA 685-784)

3 Images



#### Overview

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | N-Cadherin (CDH2)  |
| Binding Specificity: | AA 685-784   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This N-Cadherin antibody is un-conjugated  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

## **Product Details**


| Purpose:          | Polyclonal Antibody to N-cadherin (NCAD)   |
|-------------------|--|
| Immunogen:        | Recombinant N-cadherin (NCAD) corresdonding to Ser685~Asp784 with N-terminal His Tag   |
| Isotype:          | IgG  |
| Specificity:      | The antibody is a rabbit polyclonal antibody raised against NCAD. It has been selected for its ability to recognize NCAD in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Mouse, Rat   |
| Purification:     | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |

# **Target Details**

Expiry Date:


24 months

| rarget Details      |  |
|---------------------|--|
| Target:             | N-Cadherin (CDH2)  |
| Alternative Name:   | N-cadherin (CDH2 Products)   |
| Background:         | CDw325, CD325, CDHN, CDH2, Cadherin 2 Type 1, N-Cadherin   |
| Pathways:           | Regulation of Muscle Cell Differentiation, Cell-Cell Junction Organization, Synaptic Membrane    |
| Application Details |  |
| Application Notes:  | Western blotting: 0.2-2 μg/mL  |
|                     | 1:250-2500 Immunohistochemistry: 5-20 µg/mL  |
|                     | 1:25-100 Immunocytochemistry: 5-20 μg/mL   |
|                     | 1:25-100 Optimal working dilutions must be determined by end user.                               |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
|                     | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious          |
|                     | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
|                     | date under appropriate storage condition.  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Buffer:             | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.                                 |
| Preservative:       | ProClin  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be               |
|                     | handled by trained staff only.   |
| Storage:            | 4 °C,-20 °C  |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without  |
|                     | detectable loss of activity. Avoid repeated freeze-thaw cycles.                                  |
| Francisco Deter     | O.A. v. sueth s  |




## **Immunohistochemistry**

**Image 1.** Detection of NCAD in Human Glioma Tissue using Polyclonal Antibody to N-cadherin (NCAD)



#### **Immunohistochemistry**

**Image 2.** Detection of NCAD in Human Breast cancer Tissue using Polyclonal Antibody to N-cadherin (NCAD)



# **Immunohistochemistry**

**Image 3.** Detection of NCAD in Human Kidney Tissue using Polyclonal Antibody to N-cadherin (NCAD)