antibodies -online.com





anti-TNNC2 antibody (AA 2-160)

2 Images



Overview

Quantity:	100 μL
Target:	TNNC2
Binding Specificity:	AA 2-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNNC2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Troponin C Type 2, Fast (TNNC2)
Immunogen:	Recombinant Troponin C Type 2, Fast (TNNC2) corresdonding to Thr2~Gln160 with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TNNC2. It has been selected for its ability to recognize TNNC2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Storage Comment:


Expiry Date:


Target Details	
Target:	TNNC2
Alternative Name:	Troponin C Type 2, Fast (TNNC2 Products)
Background:	Troponin C, skeletal muscle
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	1:200-800 Immunohistochemistry: 5-20 μg/mL
	1:20-80 Immunocytochemistry: 5-20 μg/mL
	1:20-80 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Otamana Oamanaanti	Others at 480 for fragrens to a Channel at 2000 in a magnitude fragt fragren for the control of the channel of

detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without





Western Blotting

Image 1. Detection of TNNC2 in Rat Muscle lysate using Polyclonal Antibody to Troponin C Type 2, Fast (TNNC2)

Western Blotting

Image 2. Detection of Recombinant TNNC2, Human using Polyclonal Antibody to Troponin C Type 2, Fast (TNNC2)