# antibodies - online.com







# anti-NOX1 antibody (AA 235-488)



# **Images**



( )	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	1 V I	ew

Quantity:	100 μL	
Target:	NOX1	
Binding Specificity:	AA 235-488	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NOX1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

## **Product Details**


Purpose:	Polyclonal Antibody to Nicotinamide Adenine Dinucleotide Phosphate Oxidase 1 (NOX1)	
Immunogen:	Recombinant Nicotinamide Adenine Dinucleotide Phosphate Oxidase 1 (NOX1) corresdonding to Met235~Asn488 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against NOX1. It has been selected for its ability to recognize NOX1 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

## **Target Details**

t: NOX1
---------


# **Target Details**

•		
Alternative Name:	Nicotinamide Adenine Dinucleotide Phosphate Oxidase 1 (NOX1 Products)	
Background:	MOX1, GP91-2, NOH1, NADPH Oxidase 1, Mitogenic Oxidase(Pyridine Nucleotide-Dependent	
	Superoxide-Generating, NADH/NADPH mitogenic oxidase subunit P65-MOX	
Pathways:	Regulation of Systemic Arterial Blood Pressure by Hormones, Proton Transport	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL Immunocytochemistry in formalin fixed cells: 5-20 μg/mL	
	Immunohistochemistry in formalin fixed frozen section: 5-20 $\mu g/mL$ Immunohistochemistry in	
	paraffin section: 5-20 $\mu g/mL$ Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	




## **Immunohistochemistry**

Image 1. Detection of NOX1 in Human Cerebrum Tissueusing Polyclonal Antibody to Nicotinamide AdenineDinucleotide Phosphate Oxidase 1 (NOX1)



#### **Western Blotting**

**Image 2.** Detection of Recombinant NOX1, Human using Polyclonal Antibody to Nicotinamide Adenine Dinucleotide Phosphate Oxidase 1 (NOX1)



## **Immunohistochemistry**

**Image 3.** Detection of NOX1 in Human Liver Tissue using Polyclonal Antibody to Nicotinamide Adenine Dinucleotide Phosphate Oxidase 1 (NOX1)

Please check the product details page for more images. Overall 7 images are available for ABIN7439315.