antibodies -online.com





anti-RORC antibody (AA 212-461)





Go to Product page

\sim				
()\	10	K\ / I	01	A /
1 1	ve.	I \/ I	-1	/\/


Quantity:	100 μL
Target:	RORC
Binding Specificity:	AA 212-461
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RORC antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Purpose:	Polyclonal Antibody to RAR Related Orphan Receptor Gamma (RORg)	
Immunogen:	Recombinant RAR Related Orphan Receptor Gamma (RORg) corresdonding to Glu212~Ser461 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against RORg. It has been selected for its ability to recognize RORg in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


Target Details

RORC		
RAR Related Orphan Receptor Gamma (RORC Products)		
RORC, NR1F3, RORG, RZRG, TOR, Nuclear receptor subfamily 1 group F member 3, RAR-related		
orphan receptor C, Retinoid-related orphan receptor-gamma		
Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway		
Western blotting: 0.5-2 μg/mL		
1:500-2000 Immunohistochemistry: 5-20 µg/mL		
1:50-200 Immunocytochemistry: 5-20 μg/mL		
1:50-200 Optimal working dilutions must be determined by end user.		
The thermal stability is described by the loss rate. The loss rate was determined by accelerated		
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious		
degradation and precipitation were observed. The loss rate is less than 5% within the expiration		
date under appropriate storage condition.		
For Research Use only		
Liquid		
0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.		
ProClin		
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be		
handled by trained staff only.		
4 °C,-20 °C		
Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without		
detectable loss of activity. Avoid repeated freeze-thaw cycles.		
24 months		




Immunohistochemistry

Image 1. Detection of RORg in Human Liver cancer Tissue using Polyclonal Antibody to RAR Related Orphan Receptor Gamma (RORg)



Western Blotting

Image 2. Detection of RORg in Mouse Liver Tissue using Polyclonal Antibody to RAR Related Orphan Receptor Gamma (RORg)



Immunohistochemistry

Image 3. Detection of RORg in Human Lung Cancer Tissue using Polyclonal Antibody to RAR Related Orphan Receptor Gamma (RORg)

Please check the product details page for more images. Overall 5 images are available for ABIN7439402.