antibodies - online.com







anti-CCL6 antibody (AA 22-116)





| \sim | | | |
|--------|-----------------------|------|---------------|
| | $ \backslash / \cap$ | r\/I | $\triangle V$ |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | CCL6 | |
| Binding Specificity: | AA 22-116 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This CCL6 antibody is un-conjugated | |
| Application: | Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC), Immunoprecipitation (IP) | |

Product Details

| Purpose: | Polyclonal Antibody to Chemokine (C-C motif) ligand 6 (CCL6) | |
|---------------|--|--|
| Immunogen: | Recombinant Chemokine (C-C motif) ligand 6 (CCL6) corresdonding to Gly22~Ala116 (Accession # P27784) | |
| Isotype: | IgG | |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against CCL6. It has been selected for its ability to recognize CCL6 in immunohistochemical staining and western blotting. | |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography | |

Target Details


Target Details

| Chemokine (C-C motif) ligand 6 (CCL6 Products) | |
|--|--|
| MRP1, SCYA6, C10, Macrophage Inflammatory Protein Related Protein 1, Small Inducible Cytokine A6 | |
| | |
| | |
| Western blotting: 0.5-2 μg/mL | |
| Immunohistochemistry: 5-20 μg/mL | |
| Immunocytochemistry: 5-20 μg/mL | |
| Optimal working dilutions must be determined by end user. | |
| The thermal stability is described by the loss rate. The loss rate was determined by accelerated | |
| thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious | |
| degradation and precipitation were observed. The loss rate is less than 5% within the expiration | |
| date under appropriate storage condition. | |
| For Research Use only | |
| | |
| Liquid | |
| PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. | |
| Sodium azide | |
| This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which | |
| should be handled by trained staff only. | |
| 4 °C,-20 °C | |
| Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without | |
| detectable loss of activity. Avoid repeated freeze-thaw cycles. | |
| 24 months | |
| | |



Western Blotting

Image 1. Detection of CCL6 in Mouse Pancreas lysate using Polyclonal Antibody to Chemokine (C-C motif) ligand 6 (CCL6)



Western Blotting

Image 2. Detection of Recombinant MRP1, Mouse using Polyclonal Antibody to Chemokine (C-C motif) ligand 6 (CCL6)