antibodies - online.com







anti-Oncostatin M Receptor antibody (AA 503-749)





		oo to i roddot page

\cup	٧	е	rv	16	5/	Λ
--------	---	---	----	----	----	---

Quantity:	100 μL
Target:	Oncostatin M Receptor (OSMR)
Binding Specificity:	AA 503-749
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Oncostatin M Receptor antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Oncostatin M Receptor (OSMR)
Immunogen:	Recombinant Oncostatin M Receptor (OSMR) corresdonding to His503~Gly749 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against OSMR. It has been selected for its ability to recognize OSMR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target: Oncostatin M Receptor (OSMR)

Target Details

Expiry Date:


24 months

Oncostatin M Receptor (OSMR Products)
OSMRB, IL-31RB, Oncostatin-M-specific receptor subunit beta, Interleukin-31 receptor subunit beta
JAK-STAT Signaling, Growth Factor Binding
Western blotting: 0.5-2 μ g/mL Immunohistochemistry: 5-20 μ g/mL Immunocytochemistry: 5-20 μ g/mL Optimal working dilutions must be determined by end user.
The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
For Research Use only
Liquid
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Sodium azide
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
4 °C,-20 °C
Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.




Western Blotting

Image 1. Detection of OSMR in Rat Serum using Polyclonal Antibody to Oncostatin M Receptor (OSMR)



Western Blotting

Image 2. Detection of Recombinant OSMR, Rat using Polyclonal Antibody to Oncostatin M Receptor (OSMR)



Immunohistochemistry

Image 3. Detection of OSMR in Rat Liver Tissue using Polyclonal Antibody to Oncostatin M Receptor (OSMR)

Please check the product details page for more images. Overall 4 images are available for ABIN7439711.