antibodies -online.com





anti-TAF1 antibody (AA 1379-1636)

2 Images



Go to Product page

Overview


Quantity:	100 μL
Target:	TAF1
Binding Specificity:	AA 1379-1636
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAF1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA, Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to TATA Box Binding Protein Associated Factor 1 (TAF1)
Immunogen:	Recombinant TATA Box Binding Protein Associated Factor 1 (TAF1) corresdonding to Thr1379~Pro1636 (Accession # P21675)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TAF1. It has been selected for its ability to recognize TAF1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography


Target Details

9 - 1 - 1 - 1	
Target:	TAF1
Alternative Name:	TATA Box Binding Protein Associated Factor 1 (TAF1 Products)
Background:	NSCL2, OF, BA2R, CCG1, CCGS, P250, TAF2A, KAT4, DYT3/TAF1, TAF1 RNA Polymerase II,
	Dystonia 3, TBP-associated factor 250 kDa, Cell cycle gene 1 protein
Application Details	
Application Notes:	Western blotting: 1-5 μg/mL Immunocytochemistry in formalin fixed cells: 5-20 μg/mL
	Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in
	paraffin section: 5-20 μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optimal
	working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Immunohistochemistry

Image 1. Detection of TAF1 in Human Kidney Tissue using Polyclonal Antibody to TATA Box Binding Protein Associated Factor 1 (TAF1)



Western Blotting

Image 2. Detection of Recombinant TAF1, Human using Polyclonal Antibody to TATA Box Binding Protein Associated Factor 1 (TAF1)