# antibodies - online.com







# anti-PKD2 antibody (AA 621-832)

**Images** 



Go to	Prod	luct	page

_					
U	V	er	VI	е	W

Quantity:	100 μL
Target:	PKD2
Binding Specificity:	AA 621-832
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKD2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))


# **Product Details**

Purpose:	Polyclonal Antibody to Protein Kinase D2 (PKD2)
Immunogen:	Recombinant Protein Kinase D2 (PKD2) corresdonding to Glu621~Ser832 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PKD2. It has been selected for its ability to recognize PKD2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

# Target Details


Target:	PKD2
Alternative Name:	Protein Kinase D2 (PKD2 Products)
Background:	PRKD2, PRK-D2, PRKD-2, PK-D2, HSPC187, nPKC-D2, Serine/threonine-protein kinase D2
Pathways:	cAMP Metabolic Process, Maintenance of Protein Location, Negative Regulation of Transporter Activity

Pathways:	cAMP Metabolic Process, Maintenance of Protein Location, Negative Regulation of Transporter  Activity
Application Details	
Application Notes:	Western blotting: 1-5 $\mu$ g/mL Immunocytochemistry in formalin fixed cells: 5-20 $\mu$ g/mL Immunohistochemistry in formalin fixed frozen section: 5-20 $\mu$ g/mL Immunohistochemistry in paraffin section: 5-20 $\mu$ g/mL Enzyme-linked Immunosorbent Assay: 0.05-2 $\mu$ g/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




# **Immunohistochemistry**

**Image 1.** Detection of PKD2 in Human Liver Tissue using Polyclonal Antibody to Protein Kinase D2 (PKD2)



# **Western Blotting**

Image 2. Detection of Recombinant PKD2, Human using Polyclonal Antibody to Protein Kinase D2 (PKD2)



# **Immunohistochemistry**

**Image 3.** Detection of PKD2 in Human Stomach cancer Tissue using Polyclonal Antibody to Protein Kinase D2 (PKD2)