antibodies -online.com





anti-ATF6 antibody (AA 423-646)

3 Images



Go to Product page

Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | ATF6 |
| Binding Specificity: | AA 423-646 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ATF6 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| Purpose: | Polyclonal Antibody to Activating Transcription Factor 6 (ATF6) |
|-------------------|--|
| Immunogen: | Recombinant Activating Transcription Factor 6 (ATF6) corresdonding to Ala423~Asn646 with N-terminal His Tag |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against ATF6. It has been selected for its ability to recognize ATF6 in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Mouse |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

| Target Details | |
|---------------------|---|
| Target: | ATF6 |
| Alternative Name: | Activating Transcription Factor 6 (ATF6 Products) |
| Background: | Processed cyclic AMP-dependent transcription factor ATF-6 alpha |
| Pathways: | ER-Nucleus Signaling, Unfolded Protein Response |
| Application Details | |
| Application Notes: | Western blotting: 0.5-3 μg/mL |
| | Immunohistochemistry: 5-30 μg/mL |
| | Immunocytochemistry: 5-30 μg/mL |
| | Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| | date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without


4 °C,-20 °C

24 months

Storage:


Expiry Date:

Storage Comment:




Immunohistochemistry

Image 1. Detection of ATF6 in Human Cerebrum Tissue using Polyclonal Antibody to Activating Transcription Factor 6 (ATF6)



Western Blotting

Image 2. Detection of Recombinant ATF6, Human using Polyclonal Antibody to Activating Transcription Factor 6 (ATF6)



Immunohistochemistry

Image 3. Detection of ATF6 in Human Pancreas Tissue using Polyclonal Antibody to Activating Transcription Factor 6 (ATF6)