antibodies .- online.com







anti-NADPH Oxidase 4 antibody (AA 210-578)



Image



_						
()	V	ρ	rv	16	$\supset I$	Λ

Quantity:	100 μL
Target:	NADPH Oxidase 4 (NOX4)
Binding Specificity:	AA 210-578
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NADPH Oxidase 4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to NADPH Oxidase 4 (NOX4)
Immunogen:	Recombinant NADPH Oxidase 4 (NOX4) corresdonding to Gly210~Ser578 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against NOX4. It has been selected for its ability to recognize NOX4 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target: NADPH Oxidase 4 (NOX4)

Target Details


Alternative Name:	NADPH Oxidase 4 (NOX4 Products)	
Background:	KOX, KOX-1, RENOX, Nicotinamide Adenine Dinucleotide Phosphate Oxidase 4	
Pathways:	Carbohydrate Homeostasis, Smooth Muscle Cell Migration	

Application Details

Restrictions:	For Research Use only		
	date under appropriate storage condition.		
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration		
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious		
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated		
	1:50-200 Optimal working dilutions must be determined by end user.		
	1:50-200 Immunocytochemistry: 5-20 µg/mL		
	1:500-2000 Immunohistochemistry: 5-20 µg/mL		
Application Notes:	Western blotting: 0.5-2 μg/mL		

Handling

Format:	Liquid
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Western Blotting

Image 1. Detection of Recombinant NOX4, Mouse using Polyclonal Antibody to NADPH Oxidase 4 (NOX4)