# antibodies - online.com







# anti-BLK antibody (AA 241-494)



# **Images**



/ //	10	K / /	OIA.
1 11	$/ \square$	1 \/	$\square \backslash \backslash \backslash \backslash$
$\cup$	$^{\prime}$	1 V I	iew

Quantity:	100 μL	
Target:	BLK	
Binding Specificity:	AA 241-494	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BLK antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)	


#### **Product Details**

Purpose:	Polyclonal Antibody to B-Lymphoid Tyrosine Kinase (BLK)	
Immunogen:	Recombinant B-Lymphoid Tyrosine Kinase (BLK) corresdonding to Leu241~Tyr494	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against BLK. It has been selected for its ability to recognize BLK in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse, Pig, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

# Target Details


Target:	BLK	
Alternative Name:	B-Lymphoid Tyrosine Kinase (BLK Products)	
Background:	Tyrosine-protein kinase BLK, p55-Blk	
Pathways:	Positive Regulation of Peptide Hormone Secretion, CXCR4-mediated Signaling Events, Thromboxane A2 Receptor Signaling	

	Thromboxane A2 Receptor Signaling	
Application Details		
Application Notes:	Western blotting: 1-5 $\mu$ g/mL Immunocytochemistry in formalin fixed cells: 5-20 $\mu$ g/mL Immunohistochemistry in formalin fixed frozen section: 5-20 $\mu$ g/mL Immunohistochemistry in paraffin section: 5-20 $\mu$ g/mL Enzyme-linked Immunosorbent Assay: 0.05-2 $\mu$ g/mL Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	




#### **Western Blotting**

**Image 1.** Detection of BLK in Rat Lymph node lysate using Polyclonal Antibody to B-Lymphoid Tyrosine Kinase (BLK)



#### **Immunohistochemistry**

**Image 2.** Detection of BLK in Human Lymph node Tissue using Polyclonal Antibody to B-Lymphoid Tyrosine Kinase (BLK)



### **Immunohistochemistry**

**Image 3.** Detection of BLK in Human Breast cancer Tissue using Polyclonal Antibody to B-Lymphoid Tyrosine Kinase (BLK)

Please check the product details page for more images. Overall 6 images are available for ABIN7441157.