antibodies .- online.com







anti-SIRT3 antibody (AA 5-257)





Overview


Quantity:	100 μL
Target:	SIRT3
Binding Specificity:	AA 5-257
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SIRT3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Sirtuin 3 (SIRT3)
Immunogen:	Recombinant Sirtuin 3 (SIRT3) corresdonding to Gly5~Arg257 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against SIRT3. It has been selected for its ability to recognize SIRT3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	SIRT3


Target Details

Application Details Application Notes: Western blotting: 0.5-5 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 0.0 µg/mL Immunocytochemistry: 0.000 µg/mL Immunocytochemistry: 5-20 µg/mL Immunocytochemistry: 5-		
Application Details Application Notes: Western blotting: 0.5-5 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemiz 20 μg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by act thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the edate under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which shandled by trained staff only. Storage: 4 °C, -20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two years detectable loss of activity. Avoid repeated freeze-thaw cycles.	Alternative Name:	Sirtuin 3 (SIRT3 Products)
Application Notes: Western blotting: 0.5-5 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemic 20 μg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by acc thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the endate under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sinhandled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year widetectable loss of activity. Avoid repeated freeze-thaw cycles.	Background:	SIR2L3, Silent Mating Type Information Regulation 2 Homolog 3, NAD-dependent protein deacetylase sirtuin-3, mitochondrial, SIR2-like protein 3, Regulatory protein SIR2 homolog 3
20 µg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by acc thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the edate under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which stee handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year of detectable loss of activity. Avoid repeated freeze-thaw cycles.	Application Details	
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the edate under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sinandled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year and detectable loss of activity. Avoid repeated freeze-thaw cycles.	Application Notes:	Western blotting: 0.5-5 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.
Format: Liquid Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which sl handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year widetectable loss of activity. Avoid repeated freeze-thaw cycles.	Comment:	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
Format: Liquid Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which st handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year widetectable loss of activity. Avoid repeated freeze-thaw cycles.	Restrictions:	For Research Use only
Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which shandled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two years detectable loss of activity. Avoid repeated freeze-thaw cycles.	Handling	
Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which standled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year of detectable loss of activity. Avoid repeated freeze-thaw cycles.	Format:	Liquid
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which so handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year wide detectable loss of activity. Avoid repeated freeze-thaw cycles.	Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year of detectable loss of activity. Avoid repeated freeze-thaw cycles.	Preservative:	ProClin
Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year of detectable loss of activity. Avoid repeated freeze-thaw cycles.	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
detectable loss of activity. Avoid repeated freeze-thaw cycles.	Storage:	4 °C,-20 °C
Expiry Date: 24 months	Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
	Expiry Date:	24 months




Immunohistochemistry

Image 1. Detection of SIRT3 in Mouse Liver Tissue using Polyclonal Antibody to Sirtuin 3 (SIRT3)



Western Blotting

Image 2. Detection of SIRT3 in Mouse Heart Tissue using Polyclonal Antibody to Sirtuin 3 (SIRT3)



Western Blotting

Image 3. Detection of Recombinant SIRT3, Mouse using Polyclonal Antibody to Sirtuin 3 (SIRT3)

Please check the product details page for more images. Overall 4 images are available for ABIN7441238.