# antibodies -online.com







## anti-PSMD10 antibody (AA 1-226)



Image



$\sim$			
	$  \backslash / \cap$	r\/I	$\square$

Quantity:	100 μL	
Target:	PSMD10	
Binding Specificity:	AA 1-226	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PSMD10 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**


Purpose:	Polyclonal Antibody to Proteasome 26S Subunit, Non ATPase 10 (PSMD10)	
lmmunogen:	Recombinant Proteasome 26S Subunit, Non ATPase 10 (PSMD10) corresdonding to Met1~Gly226 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against PSMD10. It has been selected for its ability to recognize PSMD10 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

#### **Target Details**

Target: PSMD10
----------------

### **Target Details**

rarget Details		
Alternative Name:	Proteasome 26S Subunit, Non ATPase 10 (PSMD10 Products)	
Background:	P28, Gankyrin, 26S proteasome non-ATPase regulatory subunit 10, 26S proteasome regulatory subunit p28	
Pathways:	Mitotic G1-G1/S Phases, DNA Replication, Maintenance of Protein Location, Synthesis of DNA, Ubiquitin Proteasome Pathway	
Application Details		
Application Notes:	Western blotting: 1-5 $\mu$ g/mL Immunocytochemistry in formalin fixed cells: 5-20 $\mu$ g/mL Immunohistochemistry in formalin fixed frozen section: 5-20 $\mu$ g/mL Immunohistochemistry in paraffin section: 5-20 $\mu$ g/mL Enzyme-linked Immunosorbent Assay: 0.05-2 $\mu$ g/mL Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



#### **Western Blotting**

**Image 1.** Detection of Recombinant PSMD10, Human using Polyclonal Antibody to Proteasome 26S Subunit, Non ATPase 10 (PSMD10)