antibodies -online.com





anti-ATP5J antibody (AA 1-108)





Go to Product page

_							
0	V	е	r١	/1	е	V	1


Quantity:	100 μL	
Target:	ATP5J	
Binding Specificity:	AA 1-108	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ATP5J antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Purpose:	Polyclonal Antibody to Coupling Factor 6 (CF6)	
Immunogen:	Recombinant Coupling Factor 6 (CF6) corresdonding to Met1~Ala108 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against CF6. It has been selected for its ability to recognize CF6 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


Target Details

rarget Details		
Target:	ATP5J	
Alternative Name:	Coupling Factor 6 (ATP5J Products)	
Background:	ATP5J, ATP5A, ATP5, ATPM, ATP Synthase,H+ Transporting,Mitochondrial Fo Complex,Subunit	
	F6, ATP synthase-coupling factor 6, mitochondrial	
Pathways:	Proton Transport, Ribonucleoside Biosynthetic Process	
Application Details		
Application Notes:	Western blotting: 1-5 μg/mL Immunocytochemistry in formalin fixed cells: 5-20 μg/mL	
	Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in	
	paraffin section: 5-20 μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optimal	
	working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	




Immunohistochemistry

Image 1. Detection of CF6 in Human Prostate cancer Tissue using Polyclonal Antibody to Coupling Factor 6 (CF6)



Western Blotting

Image 2. Detection of Recombinant CF6, Human using Polyclonal Antibody to Coupling Factor 6 (CF6)



Immunohistochemistry

Image 3. Detection of CF6 in Human Liver cancer Tissue using Polyclonal Antibody to Coupling Factor 6 (CF6)

Please check the product details page for more images. Overall 7 images are available for ABIN7441733.