antibodies -online.com





anti-C190RF80 antibody (AA 16-198)

2 Images



Go to Product page

_					
U	V	er	VI	е	W


Quantity:	100 μL	
Target:	C190RF80	
Binding Specificity:	AA 16-198	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This C190RF80 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Angiopoietin Like Protein 8 (ANGPTL8)	
Immunogen:	Recombinant Angiopoietin Like Protein 8 (ANGPTL8) corresdonding to Val16~Ala198 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against ANGPTL8. It has been selected for its ability to recognize ANGPTL8 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


Target Details

Application Details Application Notes: Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by degradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only.	tting: 0.5-2 µg/mL tochemistry: 5-20 µg/mL tochemistry			
Background: RIFL, TD26, C19orf80, Lipasin, Betatrophin, Hepatocellular Carcinoma-Associated Application Details Application Notes: Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dillutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by thermal degradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only.	C19orf80, Lipasin, Betatrophin, Hepatocellular Carcinoma-Associated Protein TD26 tting: 0.5-2 µg/mL tochemistry: 5-20 µg/mL tochemistry: 5-20 µg/mL tking dilutions must be determined by end user. I stability is described by the loss rate. The loss rate was determined by accelerated radation test, that is, incubate the protein at 37°C for 48h, and no obvious and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. Ith Use only I containing 0.02 % Sodium azide, 50 % glycerol. Ide It contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only.	Target:	C190RF80	
Application Details Application Notes: Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by degradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only. Storage: 4 °C,-20 °C	titing: 0.5-2 µg/mL tochemistry: 5-20 µg/mL cochemistry: 5-20 µg/mL cochemistr	Alternative Name:	Angiopoietin Like Protein 8 (C190RF80 Products)	
Application Notes: Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by degradation test, that is, incubate the protein at 37°C for 48h, and no obvidegradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only. Storage: 4 °C,-20 °C	tochemistry: 5-20 µg/mL chemistry: 5-20 µg/mL ching dilutions must be determined by end user. It stability is described by the loss rate. The loss rate was determined by accelerated radation test, that is, incubate the protein at 37°C for 48h, and no obvious and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. In the Use only It containing 0.02 % Sodium azide, 50 % glycerol. It contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only.	Background:	RIFL, TD26, C19orf80, Lipasin, Betatrophin, Hepatocellular Carcinoma-Associated Protein TD26	
Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by degradation test, that is, incubate the protein at 37°C for 48h, and no obvidegradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only.	tochemistry: 5-20 µg/mL chemistry: 5-20 µg/mL ching dilutions must be determined by end user. It stability is described by the loss rate. The loss rate was determined by accelerated radation test, that is, incubate the protein at 37°C for 48h, and no obvious and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. In the Use only It containing 0.02 % Sodium azide, 50 % glycerol. It contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only.	Application Details		
Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by the remainded by the loss rate was determined by the remainded by the loss rate was determined by the remainded by the loss rate was determined by the remainded by the loss rate is less than 5°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE should be handled by trained staff only. Storage: 4 °C,-20 °C	chemistry: 5-20 µg/mL Rking dilutions must be determined by end user. It stability is described by the loss rate. The loss rate was determined by accelerated tradation test, that is, incubate the protein at 37°C for 48h, and no obvious and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. The Use only It containing 0.02 % Sodium azide, 50 % glycerol. It contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only.	Application Notes:	Western blotting: 0.5-2 μg/mL	
Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by the members and thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvide degradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANIShould be handled by trained staff only. Storage: 4 °C,-20 °C	I stability is described by the loss rate. The loss rate was determined by accelerated radation test, that is, incubate the protein at 37°C for 48h, and no obvious and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. The loss rate is less than 5% within the expiration appropriate storage condition. The Use only The loss rate is less than 5% within the expiration appropriate storage condition. The Use only The loss rate is less than 5% within the expiration appropriate storage condition. The Use only The loss rate was determined by accelerated at 30°C for 48h, and no obvious and precipitation of the expiration and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. The Use only The loss rate was determined by accelerated at 30°C for 48h, and no obvious and precipitation of the expiration and precipitation were observed. The loss rate was determined by accelerated at 40°C for 48h, and no obvious and precipitation of the expiration of the expiration and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition.		Immunohistochemistry: 5-20 μg/mL	
Comment: The thermal stability is described by the loss rate. The loss rate was determined by the loss rate was determined by the loss rate is less than 5°C for 48h, and no obvide degradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only. Storage: 4 °C,-20 °C	stability is described by the loss rate. The loss rate was determined by accelerated radation test, that is, incubate the protein at 37°C for 48h, and no obvious and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. The Use only In containing 0.02 % Sodium azide, 50 % glycerol. It contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only.		Immunocytochemistry: 5-20 μg/mL	
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvidegradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANG should be handled by trained staff only. Storage: 4 °C,-20 °C	radation test, that is, incubate the protein at 37°C for 48h, and no obvious and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. The Use only The Containing 0.02 % Sodium azide, 50 % glycerol. The Containing 0.02 % Sodium azide, 50 % glycerol. The Containing 0.02 % Sodium azide, 50 % glycerol. The Containing 0.02 % Sodium azide a POISONOUS AND HAZARDOUS SUBSTANCE which and led by trained staff only.		Optimal working dilutions must be determined by end user.	
degradation and precipitation were observed. The loss rate is less than 5% within date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTAN should be handled by trained staff only. Storage: 4 °C,-20 °C	and precipitation were observed. The loss rate is less than 5% within the expiration appropriate storage condition. th Use only de containing 0.02 % Sodium azide, 50 % glycerol. de t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only.	Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANG should be handled by trained staff only. Storage: 4 °C,-20 °C	appropriate storage condition. th Use only containing 0.02 % Sodium azide, 50 % glycerol. de t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only. for frequent use. Stored at -20°C in a manual defrost freezer for two year without		thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
Restrictions: For Research Use only Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANI should be handled by trained staff only. Storage: 4 °C,-20 °C	th Use only containing 0.02 % Sodium azide, 50 % glycerol. de t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only. for frequent use. Stored at -20°C in a manual defrost freezer for two year without		degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
Handling Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANG should be handled by trained staff only. Storage: 4 °C,-20 °C	t containing 0.02 % Sodium azide, 50 % glycerol. de t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only.		date under appropriate storage condition.	
Format: Liquid Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANG should be handled by trained staff only. Storage: 4 °C,-20 °C	de t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only. for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Restrictions:	For Research Use only	
Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANGE should be handled by trained staff only. Storage: 4 °C,-20 °C	de t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only. for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Handling		
Preservative: Sodium azide Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANG should be handled by trained staff only. Storage: 4 °C,-20 °C	t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only. for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Format:	Liquid	
Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANGE should be handled by trained staff only. Storage: 4 °C,-20 °C	t contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which andled by trained staff only. for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
should be handled by trained staff only. Storage: 4 °C,-20 °C	andled by trained staff only. If for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Preservative:	Sodium azide	
Storage: 4 °C,-20 °C	for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
			should be handled by trained staff only.	
Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two		Storage:	4 °C,-20 °C	
· ·	oss of activity. Avoid repeated freeze-thaw cycles.	Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
detectable loss of activity. Avoid repeated freeze-thaw cycles.			detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date: 24 months		Expiry Date:	24 months	



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant ANGPTL8, Mouse using Polyclonal Antibody to Angiopoietin Like Protein 8 (ANGPTL8)