
# antibodies -online.com





# anti-HIF1AN antibody (AA 31-349)



**Images** 



Go to Product page

( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷


Quantity:	100 μL	
Target:	HIF1AN	
Binding Specificity:	AA 31-349	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HIF1AN antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

# **Product Details**

Purpose:	Polyclonal Antibody to Hypoxia Inducible Factor 1 Alpha Subunit Inhibitor (HIF1aN)	
lmmunogen:	Recombinant Hypoxia Inducible Factor 1 Alpha Subunit Inhibitor (HIF1aN) corresdonding to Gln31~Asn349 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against HIF1aN. It has been selected for its ability to recognize HIF1aN in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse, Pig, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


# **Target Details**

Target:	HIF1AN	
Alternative Name:	Hypoxia Inducible Factor 1 Alpha Subunit Inhibitor (HIF1AN Products)	
Background:	FIH1, Peptide-Aspartate Beta-Dioxygenase, Hypoxia-inducible factor asparagine hydroxylase	
Pathways:	Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL	
	Immunohistochemistry: 5-20 μg/mL	
	Immunocytochemistry: 5-20 μg/mL	
	Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	




# **Immunohistochemistry**

**Image 1.** Detection of HIF1aN in Human Prostate Tissue using Polyclonal Antibody to Hypoxia Inducible Factor 1 Alpha Subunit Inhibitor (HIF1aN)



### **Immunohistochemistry**

Image 2. Detection of HIF1aN in Human Hela Cells using Polyclonal Antibody to Hypoxia Inducible Factor 1 Alpha Subunit Inhibitor (HIF1aN)



# **Western Blotting**

**Image 3.** Detection of Recombinant HIF1aN, Human using Polyclonal Antibody to Hypoxia Inducible Factor 1 Alpha Subunit Inhibitor (HIF1aN)

Please check the product details page for more images. Overall 6 images are available for ABIN7441979.