antibodies -online.com





anti-EIF6 antibody (AA 622-848)



Images



Go to Product page

()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢


Quantity:	100 μL	
Target:	EIF6	
Binding Specificity:	AA 622-848	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EIF6 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Eukaryotic Translation Initiation Factor 3A (EIF3A)	
Immunogen:	Recombinant Eukaryotic Translation Initiation Factor 3A (EIF3A) corresdonding to Glu622~Val848 (Accession # P23116)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against EIF3A. It has been selected for its ability to recognize EIF3A in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Human, Mouse	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


Target Details

EIF6		
Eukaryotic Translation Initiation Factor 3A (EIF6 Products)		
EIF3S10, EIF3-A, EIF3, P167, EIF3-P170, EIF3-Theta, P180, P185, Eukaryotic translation initiation factor 3 subunit 10		
Ribonucleoprotein Complex Subunit Organization, Ribosome Assembly		
Western blotting: 0.5-5 μ g/mL Immunohistochemistry: 5-20 μ g/mL Immunocytochemistry: 5-20 μ g/mL Optimal working dilutions must be determined by end user.		
The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.		
For Research Use only		
Liquid		
0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.		
ProClin		
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.		
4 °C,-20 °C		
Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.		
,		




Immunohistochemistry

Image 1. Detection of EIF3A in Mouse Kidney Tissue using Polyclonal Antibody to Eukaryotic Translation Initiation Factor 3A (EIF3A)



Western Blotting

Image 2. Detection of Recombinant EIF3A, Mouse using Polyclonal Antibody to Eukaryotic Translation Initiation Factor 3A (EIF3A)



Immunohistochemistry

Image 3. Detection of EIF3A in Mouse Brain Tissue using Polyclonal Antibody to Eukaryotic Translation Initiation Factor 3A (EIF3A)