
# antibodies - online.com







## anti-CXCL9 antibody (AA 23-125)



Image



( )	ve	K\ /		
	$\cup$	1 V/	Щ.	V۷


Quantity:	100 μL	
Target:	CXCL9	
Binding Specificity:	AA 23-125	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CXCL9 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

#### **Product Details**

Purpose:	Polyclonal Antibody to Monokine Induced By Interferon Gamma (MIg)	
Immunogen:	Recombinant Monokine Induced By Interferon Gamma (MIg) corresdonding to Thr23~Thr125 with N-terminal His and GST Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against Mlg. It has been selected for its ability to recognize Mlg in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Pig	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

### **Target Details**

CXCL9	
Monokine Induced By Interferon Gamma (CXCL9 Products)	
CXCL9, CMK, Humig, MI-G, SCYB9, Crg-10, Chemokine (C-X-C motif) ligand 9, Small-inducible	
cytokine B9, Gamma-interferon-induced monokine	
Western blotting: 0.5-2 μg/mL	
Immunohistochemistry: 5-20 μg/mL	
Immunocytochemistry: 5-20 μg/mL	
Optimal working dilutions must be determined by end user.	
The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
date under appropriate storage condition.	
For Research Use only	
Liquid	
0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
handled by trained staff only.	
4 °C,-20 °C	
Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
detectable loss of activity. Avoid repeated freeze-thaw cycles.	
24 months	



#### **Western Blotting**

**Image 1.** Detection of Recombinant MIg, Human using Polyclonal Antibody to Monokine Induced By Interferon Gamma (MIg)