# antibodies - online.com





Datasheet for ABIN7442504

## anti-TIE1 antibody (AA 435-609)






_							
0	V	е	r١	/1	е	V	1

Quantity:	100 μL
Target:	TIE1
Binding Specificity:	AA 435-609
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TIE1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Product Details  Purpose:	Polyclonal Antibody to Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1)
Purpose:	Protein 1 (Tie1)  Recombinant Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1
Purpose:  Immunogen:	Protein 1 (Tie1)  Recombinant Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1  (Tie1) corresdonding to Asp435~Gly609 with N-terminal His Tag
Purpose:  Immunogen:  Isotype:	Protein 1 (Tie1)  Recombinant Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1) corresdonding to Asp435~Gly609 with N-terminal His Tag  IgG  The antibody is a rabbit polyclonal antibody raised against Tie1. It has been selected for its

### **Target Details**


Target:	TIE1
Alternative Name:	Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (TIE1 Products)
Background:	JTK14, Tyrosine Kinase With Immunoglobulin And Epidermal Growth Factor Homology Domains 1, Tyrosine-protein kinase receptor Tie-1
Pathways:	RTK Signaling
Application Details	

Application Notes:	Western blotting: 1-5 μg/mL Immunocytochemistry in formalin fixed cells: 5-20 μg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 μg/mL Immunohistochemistry in paraffin section: 5-20 μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



#### Immunohistochemistry

**Image 1.** Detection of Tie1 in Human Kidney Tissue using Polyclonal Antibody to Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1)



#### **Western Blotting**

**Image 2.** Detection of Tie1 in Human K562 Cells using Polyclonal Antibody to Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1)

#### **Western Blotting**

**Image 3.** Detection of Recombinant Tie1, Human using Polyclonal Antibody to Tyrosine Kinase With Immunoglobulin Like And EGF Like Domains Protein 1 (Tie1)

Please check the product details page for more images. Overall 4 images are available for ABIN7442504.