
antibodies -online.com









Overview

Quantity:	100 μL
Target:	HIPK2
Binding Specificity:	AA 977-1198
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIPK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Homeodomain Interacting Protein Kinase 2 (HIPK2)
Immunogen:	Recombinant Homeodomain Interacting Protein Kinase 2 (HIPK2) corresdonding to Ser977~IIe1198
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HIPK2. It has been selected for its ability to recognize HIPK2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	HIPK2
Alternative Name:	Homeodomain Interacting Protein Kinase 2 (HIPK2 Products)
Pathways:	Cell Division Cycle

Application Details

Application Notes:	Western blotting: 1-5 μg/mL Immunocytochemistry in formalin fixed cells: 5-20 μg/mL
	Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in
	paraffin section: 5-20 μg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μg/mL Optimal
	working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant HIPK2, Human using Polyclonal Antibody to Homeodomain Interacting Protein Kinase 2 (HIPK2)