

Datasheet for ABIN7442618  
**anti-CD40 Ligand antibody (AA 1-261)**

## 6 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	CD40 Ligand (CD40LG)
Binding Specificity:	AA 1-261
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD40 Ligand antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## Product Details

Purpose:	Polyclonal Antibody to Cluster Of Differentiation 40 Ligand (CD40L)
Immunogen:	Recombinant Cluster Of Differentiation 40 Ligand (CD40L) corresponding to Met1~Leu261 (Accession # P29965) with N-terminal His and GST Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CD40L. It has been selected for its ability to recognize CD40L in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details


Target:	CD40 Ligand (CD40LG)
Alternative Name:	CD40 Ligand ( <a href="#">CD40LG Products</a> )
Background:	CD154, TRAP, HIGM1, IGM, IMD3, TBAM, T-BAM, TNFSF5, Gp39, TNF Superfamily Member 5, Hyper-IgM Syndrome, TNF-Related Activation Protein, T-Cell B-Cell Activating Molecule
Pathways:	<a href="#">NF-kappaB Signaling</a> , <a href="#">Production of Molecular Mediator of Immune Response</a> , <a href="#">Cancer Immune Checkpoints</a>

## Application Details

Application Notes:	Western blotting: 1-5 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




#### Immunohistochemistry

**Image 1.** Detection of CD40L in Human Liver Tissue using Polyclonal Antibody to Cluster Of Differentiation 40 Ligand (CD40L)



#### Western Blotting

**Image 2.** Detection of Recombinant CD40L, Human using Polyclonal Antibody to Cluster Of Differentiation 40 Ligand (CD40L)



#### Immunohistochemistry

**Image 3.** Detection of CD40L in Human Cardiac Muscle Tissue using Polyclonal Antibody to Cluster Of Differentiation 40 Ligand (CD40L)

Please check the [product details page](#) for more images. Overall 6 images are available for ABIN7442618.