# antibodies -online.com





## anti-Asialoglycoprotein Receptor 1 antibody (AA 80-281)





Go to Product page

| $\sim$ |     |        |               |     |            |   |    |
|--------|-----|--------|---------------|-----|------------|---|----|
|        | 1// | $\Box$ | $r \setminus$ | / [ | $\bigcirc$ | 1 | ٨, |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | Asialoglycoprotein Receptor 1 (ASGR1)  |  |
| Binding Specificity: | AA 80-281  |  |
| Reactivity:          | Mouse  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This Asialoglycoprotein Receptor 1 antibody is un-conjugated   |  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |  |

#### **Product Details**


| Purpose:      | Polyclonal Antibody to Asialoglycoprotein Receptor 1 (ASGR1)   |  |
|---------------|--|--|
| Immunogen:    | Recombinant Asialoglycoprotein Receptor 1 (ASGR1) corresdonding to Thr80~Asp281 with N-terminal His Tag  |  |
| Isotype:      | IgG  |  |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against ASGR1. It has been selected for its ability to recognize ASGR1 in immunohistochemical staining and western blotting. |  |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography   |  |

### **Target Details**

| Target: Asialoglycoprotein Receptor 1 (ASGR1) |
|---|
|---|


#### **Target Details**

| l arget Details     |  |  |
|---------------------|--|--|
| Alternative Name:   | Asialoglycoprotein Receptor 1 (ASGR1 Products)   |  |
| Background:         | CLEC4H1, HL-1, ASGPR 1, Asialoglyco protein Receptor, C-type lectin domain family 4 member       |  |
|                     | H1, Hepatic lectin H1  |  |
| Pathways:           | Thyroid Hormone Synthesis  |  |
| Application Details |  |  |
| Application Notes:  | Western blotting: 0.5-2 μg/mL  |  |
|                     | Immunohistochemistry: 5-20 μg/mL   |  |
|                     | Immunocytochemistry: 5-20 μg/mL  |  |
|                     | Optimal working dilutions must be determined by end user.  |  |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |  |
|                     | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious          |  |
|                     | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |  |
|                     | date under appropriate storage condition.  |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Buffer:             | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.                                 |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be               |  |
|                     | handled by trained staff only.   |  |
| Storage:            | 4 °C,-20 °C  |  |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without  |  |
|                     | detectable loss of activity. Avoid repeated freeze-thaw cycles.                                  |  |
| Expiry Date:        | 24 months  |  |




#### **Immunohistochemistry**

**Image 1.** Detection of ASGR1 in Mouse Liver Tissue using Polyclonal Antibody to Asialoglycoprotein Receptor 1 (ASGR1)



#### **Western Blotting**

Image 2. Detection of Recombinant ASGR1, Mouse usingPolyclonal Antibody to Asialoglycoprotein Receptor 1(ASGR1)



#### **Immunohistochemistry**

**Image 3.** Detection of ASGR1 in Mouse Stomach Tissue using Polyclonal Antibody to Asialoglycoprotein Receptor 1 (ASGR1)