
antibodies -online.com





anti-STAM antibody (AA 199-464)

2 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	STAM
Binding Specificity:	AA 199-464
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAM antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Signal Transducing Adaptor Molecule 1 (STAM1)
Immunogen:	Recombinant Signal Transducing Adaptor Molecule 1 (STAM1) corresdonding to Pro199~Met464 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against STAM1. It has been selected for its ability to recognize STAM1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	STAM
Alternative Name:	Signal Transducing Adaptor Molecule 1 (STAM Products)
Background:	SH3 Domain And ITAM Motif
Pathways:	EGFR Signaling Pathway, EGFR Downregulation
Application Details	


Application Notes:

	20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Western blotting: $0.5-5 \mu g/mL$ Immunohistochemistry: $5-20 \mu g/mL$ Immunocytochemistry: $5-20 \mu g/mL$


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant STAM1, Mouse using Polyclonal Antibody to Signal Transducing Adaptor Molecule 1 (STAM1)