-online.com antibodies

## Datasheet for ABIN7443026 anti-HMGN2 antibody (AA 2-81)

5 Images



#### Overview


Quantity:	100 µL
Target:	HMGN2
Binding Specificity:	AA 2-81
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGN2 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

### Product Details

Purpose:	Polyclonal Antibody to High Mobility Group Nucleosome Binding Domain Containing Protein 2 (HMGN2)
Immunogen:	Recombinant High Mobility Group Nucleosome Binding Domain Containing Protein 2 (HMGN2) corresdonding to Pro2~Gln81 with N-terminal His Tag
lsotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HMGN2. It has been selected for its ability to recognize HMGN2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7443026 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Target Details		
Target:	HMGN2	
Alternative Name:	High Mobility Group Nucleosome Binding Domain Containing Protein 2 (HMGN2 Products)	
Background:	HMG17, High Mobility Group Protein 17, Non-histone chromosomal protein HMG-17	
Application Details		
Application Notes:	Western blotting: 0.5-2 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	



Western Blot

←17kDa

kDa 70

44

33

26 22 18


14 10

#### Immunohistochemistry

**Image 1.** Detection of HMGN2 in Human Skin cancer Tissue using Polyclonal Antibody to High Mobility Group Nucleosome Binding Domain Containing Protein 2 (HMGN2)

# Western Blotting

**Image 2.** Detection of Recombinant HMG17, Human using Polyclonal Antibody to High Mobility Group Nucleosome Binding Domain Containing Protein 2 (HMGN2)



#### Immunohistochemistry

**Image 3.** Detection of HMGN2 in Human Stomach Tissue using Polyclonal Antibody to High Mobility Group Nucleosome Binding Domain Containing Protein 2 (HMGN2)

Please check the product details page for more images. Overall 5 images are available for ABIN7443026.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7443026 | 09/09/2023 | Copyright antibodies-online. All rights reserved.