antibodies -online.com





anti-SCO2 antibody (AA 42-266)





Go to Product page

\sim			
	IV/E	۱/۱۲	$I \cap V$

Quantity:	100 μL	
Target:	SCO2	
Binding Specificity:	AA 42-266	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SCO2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to SCO Cytochrome Oxidase Deficient Homolog 2 (SCO2)
Immunogen:	Recombinant SCO Cytochrome Oxidase Deficient Homolog 2 (SCO2) corresdonding to Gly42~Ser266 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against SCO2. It has been selected for its ability to recognize SCO2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target Details


Alternative Name:	SCO Cytochrome Oxidase Deficient Homolog 2 (SCO2 Products)
Background:	SC01L
Pathways:	Transition Metal Ion Homeostasis, Warburg Effect

Application Details

Application Notes:	Western blotting: 0.5-5 μ g/mL Immunohistochemistry: 5-20 μ g/mL Immunocytochemistry: 5-
	20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Immunohistochemistry

Image 1. Detection of SCO2 in Human Stomach Tissue using Polyclonal Antibody to SCO Cytochrome Oxidase Deficient Homolog 2 (SCO2)



Western Blotting

Image 2. Detection of Recombinant SCO2, Human using Polyclonal Antibody to SCO Cytochrome Oxidase Deficient Homolog 2 (SCO2)