

Datasheet for ABIN7443464  
**anti-TANK antibody (AA 131-369)**

## 3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	TANK
Binding Specificity:	AA 131-369
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TANK antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## Product Details

Purpose:	Polyclonal Antibody to TRAF Family Member Associated NFKB Activator (TANK)
Immunogen:	Recombinant TRAF Family Member Associated NFKB Activator (TANK) corresponding to Leu131~Ile369
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TANK. It has been selected for its ability to recognize TANK in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## Target Details

Target:	TANK
Alternative Name:	TRAF Family Member Associated NFkB Activator ( <a href="#">TANK Products</a> )
Background:	TRAF2, I-TRAF, TRAF-interacting protein
Pathways:	<a href="#">p53 Signaling</a> , <a href="#">TLR Signaling</a> , <a href="#">Activation of Innate immune Response</a>

## Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL 1:500-2000 Immunohistochemistry: 5-20 µg/mL 1:50-200 Immunocytochemistry: 5-20 µg/mL 1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




#### Western Blotting

**Image 1.** Detection of TANK in Porcine Colon lysate using Polyclonal Antibody to TRAF Family Member Associated NFKB Activator (TANK)



#### Western Blotting

**Image 2.** Detection of Recombinant TANK, Human using Polyclonal Antibody to TRAF Family Member Associated NFKB Activator (TANK)



#### Immunohistochemistry

**Image 3.** Detection of TANK in Human Kidney Tissue using Polyclonal Antibody to TRAF Family Member Associated NFKB Activator (TANK)