# antibodies - online.com







## anti-CD74 antibody (AA 64-205)



## **Images**



| 0 | 1 / | -        | K   | /1  | -        | 1 A |
|---|-----|----------|-----|-----|----------|-----|
| u | \/  | $\vdash$ | ı ۱ | / I | $\vdash$ | 1/1 |
|   |     |          |     |     |          |     |


| Quantity:            | 100 μL                                                                                                 |
|----------------------|--------------------------------------------------------------------------------------------------------|
| Target:              | CD74                                                                                                   |
| Binding Specificity: | AA 64-205                                                                                              |
| Reactivity:          | Mouse                                                                                                  |
| Host:                | Rabbit                                                                                                 |
| Clonality:           | Polyclonal                                                                                             |
| Conjugate:           | This CD74 antibody is un-conjugated                                                                    |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

## **Product Details**

| Purpose:          | Polyclonal Antibody to Cluster Of Differentiation 74 (CD74)                                                                                                                    |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunogen:        | Recombinant Cluster Of Differentiation 74 (CD74) corresdonding to Leu64~Val205                                                                                                 |
| Isotype:          | IgG                                                                                                                                                                            |
| Specificity:      | The antibody is a rabbit polyclonal antibody raised against CD74. It has been selected for its ability to recognize CD74 in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Rat                                                                                                                                                                            |
| Purification:     | Antigen-specific affinity chromatography followed by Protein A affinity chromatography                                                                                         |


## **Target Details**

| Target:             | CD74                                                                                                                                                                                                                                                                                                                                |  |  |
|---------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|
| Alternative Name:   | CD74 (CD74 Products)                                                                                                                                                                                                                                                                                                                |  |  |
| Background:         | MHCDG, HLA-DG, HLADG, DHLAG, Ia-GAMMA, HLA class II histocompatibility antigen gamma chain, Ia antigen-associated invariant chain, HLA-DR antigens-associated invariant chain                                                                                                                                                       |  |  |
| Pathways:           | Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Negative Regulation of intrinsic apoptotic Signaling, Cancer Immune Checkpoints                                                                                                                                                |  |  |
| Application Details |                                                                                                                                                                                                                                                                                                                                     |  |  |
| Application Notes:  | Western blotting: $0.5-2~\mu g/mL$ Immunocytochemistry in formalin fixed cells: $5-20~\mu g/mL$ Immunohistochemistry in formalin fixed frozen section: $5-20~\mu g/mL$ Immunohistochemistry in paraffin section: $5-20~\mu g/mL$ Optimal working dilutions must be determined by end user.                                          |  |  |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |  |  |
| Restrictions:       | For Research Use only                                                                                                                                                                                                                                                                                                               |  |  |
| Handling            |                                                                                                                                                                                                                                                                                                                                     |  |  |
| Format:             | Liquid                                                                                                                                                                                                                                                                                                                              |  |  |
| Buffer:             | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.                                                                                                                                                                                                                                                                    |  |  |
| Preservative:       | ProClin                                                                                                                                                                                                                                                                                                                             |  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                                                                                                                                                                                                                   |  |  |
| Storage:            | 4 °C,-20 °C                                                                                                                                                                                                                                                                                                                         |  |  |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.                                                                                                                                                                     |  |  |
| Expiry Date:        | 24 months                                                                                                                                                                                                                                                                                                                           |  |  |




## **Immunohistochemistry**

**Image 1.** Detection of CD74 in Mouse Ovary Tissue using Polyclonal Antibody to Cluster Of Differentiation 74 (CD74)



#### **Western Blotting**

Image 2. Detection of Recombinant MHCDG, Human using Polyclonal Antibody to Cluster Of Differentiation 74 (CD74)



## **Immunohistochemistry**

**Image 3.** Detection of CD74 in Mouse Skeletal muscle Tissue using Polyclonal Antibody to Cluster Of Differentiation 74 (CD74)

Please check the product details page for more images. Overall 7 images are available for ABIN7443722.