antibodies - online.com







anti-VEGFC antibody (AA 112-227)



Image



Overview

Quantity:	100 μL
Target:	VEGFC
Binding Specificity:	AA 112-227
Reactivity:	Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VEGFC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)


Product Details

Purpose:	Polyclonal Antibody to Vascular Endothelial Growth Factor C (VEGFC)
Immunogen:	Recombinant Vascular Endothelial Growth Factor C (VEGFC) corresdonding to Ala112~Arg227
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against VEGFC. It has been selected for its ability to recognize VEGFC in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details	
Target:	VEGFC
Alternative Name:	Vascular Endothelial Growth Factor C (VEGFC Products)
Background:	VEGF-C, Flt4-L, VRPL, Vascular Endothelial Growth Factor-Related Protein
Pathways:	RTK Signaling, Signaling Events mediated by VEGFR1 and VEGFR2
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	1:500-2000 Immunohistochemistry: 5-20 μg/mL
	1:50-200 Immunocytochemistry: 5-20 µg/mL
	1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Llandling	

Handling

The state of the s	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Western Blotting

Image 1. Detection of Recombinant VEGFC, Canine using Polyclonal Antibody to Vascular Endothelial Growth Factor C (VEGFC)