antibodies -online.com







anti-Resistin antibody (AA 22-109)



Image



| \sim | |
|---------|-------|
| ()\/\ | rview |
| \circ | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | Resistin (RETN) |
| Binding Specificity: | AA 22-109 |
| Reactivity: | Pig |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Resistin antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details


Target:

| Purpose: | Polyclonal Antibody to Resistin (RETN) |
|----------------|--|
| Immunogen: | Recombinant Resistin (RETN) corresdonding to Cys22~Pro109 |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against RETN. It has been selected for its ability to recognize RETN in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |
| Target Details | |

Resistin (RETN)

Target Details

| Resistin (RETN Products) |
|--|
| |
| ADSF, FIZZ3, RETN1, RSTN, XCP1, Adipose tissue-specific secretory factor, C/EBP-epsilon- |
| regulated myeloid-specific secreted cysteine-rich protein, Cysteine-rich secreted FIZZ3 |
| Feeding Behaviour, Smooth Muscle Cell Migration |
| |
| Western blotting: 0.5-2 μg/mL |
| 1:140-560 Immunohistochemistry: 5-20 µg/mL |
| 1:14-56 Immunocytochemistry: 5-20 μg/mL |
| 1:14-56 Optimal working dilutions must be determined by end user. |
| The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious |
| degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| date under appropriate storage condition. |
| For Research Use only |
| |
| Liquid |
| PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Sodium azide |
| This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| should be handled by trained staff only. |
| 4 °C,-20 °C |
| Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without |
| detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| 24 months |
| |



Western Blotting

Image 1. Detection of Recombinant Resistin, Porcine using Polyclonal Antibody to Resistin (RETN)