

# Datasheet for ABIN7444046 anti-STAT3 antibody (AA 335-546)





Go to Product page

( )	ve	V /	-	1 A
	$\cup$	1 \/	-	1/1
$\sim$	' V C	1 V	ı	v v

Quantity:	100 μL
Target:	STAT3
Binding Specificity:	AA 335-546
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),
	Immunocytochemistry (ICC)
Product Details	
Purpose:	
i dipose.	Polyclonal Antibody to Signal Transducer And Activator Of Transcription 3 (STAT3)
Immunogen:	Polyclonal Antibody to Signal Transducer And Activator Of Transcription 3 (STAT3)  Recombinant Signal Transducer And Activator Of Transcription 3 (STAT3) corresdonding to
·	
·	Recombinant Signal Transducer And Activator Of Transcription 3 (STAT3) corresdonding to
Immunogen:	Recombinant Signal Transducer And Activator Of Transcription 3 (STAT3) corresdonding to Arg335~Trp546 with N-terminal His Tag
Immunogen: Isotype:	Recombinant Signal Transducer And Activator Of Transcription 3 (STAT3) corresdonding to  Arg335~Trp546 with N-terminal His Tag  IgG
Immunogen: Isotype:	Recombinant Signal Transducer And Activator Of Transcription 3 (STAT3) corresdonding to Arg335~Trp546 with N-terminal His Tag  IgG  The antibody is a rabbit polyclonal antibody raised against STAT3. It has been selected for its
Immunogen:  Isotype:  Specificity:	Recombinant Signal Transducer And Activator Of Transcription 3 (STAT3) corresdonding to Arg335~Trp546 with N-terminal His Tag  IgG  The antibody is a rabbit polyclonal antibody raised against STAT3. It has been selected for its ability to recognize STAT3 in immunohistochemical staining and western blotting.

## **Target Details**

Alternative Name:	STAT3 (STAT3 Products)	
Background:	APRF, Acute-Phase Response Factor	
Pathways:	JAK-STAT Signaling, RTK Signaling, Interferon-gamma Pathway, Neurotrophin Signaling	
	Pathway, Dopaminergic Neurogenesis, Response to Growth Hormone Stimulus, Carbohydrate	
	Homeostasis, Stem Cell Maintenance, Hepatitis C, Protein targeting to Nucleus, Feeding	
	Behaviour, CXCR4-mediated Signaling Events, Signaling of Hepatocyte Growth Factor Receptor	

## **Application Details**

Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



### **Immunohistochemistry**

**Image 1.** Detection of STAT3 in Human Kidney Tissue using Polyclonal Antibody to Signal Transducer And Activator Of Transcription 3 (STAT3)



#### **Immunohistochemistry**

**Image 2.** Detection of STAT3 in Human Liver Tissue using Polyclonal Antibody to Signal Transducer And Activator Of Transcription 3 (STAT3)



### **Immunohistochemistry**

**Image 3.** Detection of STAT3 in Human Stomach Tissue using Polyclonal Antibody to Signal Transducer And Activator Of Transcription 3 (STAT3)

Please check the product details page for more images. Overall 4 images are available for ABIN7444046.