
antibodies -online.com





anti-BAFF antibody (AA 21-169)

3 Images



Go to Product page

\sim							
	1//	\Box	$r \setminus$	/ [\bigcirc	1	٨,


Quantity:	100 μL	
Target:	BAFF (TNFSF13B)	
Binding Specificity:	AA 21-169	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BAFF antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to B-Cell Activation Factor Receptor (BAFFR)	
lmmunogen:	Recombinant B-Cell Activation Factor Receptor (BAFFR) corresdonding to Gln21~Lys169 (Accession # D3ZNT3) with N-terminal His and GST Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against BAFFR. It has been selected for its ability to recognize BAFFR in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	


Target Details

rarget Details		
Target:	BAFF (TNFSF13B)	
Alternative Name:	B-Cell Activation Factor Receptor (TNFSF13B Products)	
Background:	CD268, TNFRSF13C, TNFRSF13-C, BAFF-R, BAFFR, BR3, Tumor Necrosis Factor Receptor	
	Superfamily Member 13C	
Pathways:	NF-kappaB Signaling, Production of Molecular Mediator of Immune Response	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL	
	Immunohistochemistry: 5-20 μg/mL	
	Immunocytochemistry: 5-20 μg/mL	
	Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	
Expiry Date:	24 months	




Western Blotting

Image 1. Detection of BAFFR in Rat Spleen lysate using Polyclonal Antibody to B-Cell Activation Factor Receptor (BAFFR)



Western Blotting

Image 2. Detection of Recombinant BAFFR, Rat using Polyclonal Antibody to B-Cell Activation Factor Receptor (BAFFR)



Immunohistochemistry

Image 3. #VALUE!