antibodies -online.com





anti-STAM2 antibody (AA 1-377)





Go to Product page

_					
U	V	er	VI	е	W

Quantity:	100 μL	
Target:	STAM2	
Binding Specificity:	AA 1-377	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This STAM2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Signal Transducing Adaptor Molecule 2 (STAM2)	
Immunogen:	Recombinant Signal Transducing Adaptor Molecule 2 (STAM2) corresdonding to Met1~Leu377	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against STAM2. It has been selected for its ability to recognize STAM2 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Human, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details	
Target:	STAM2
Alternative Name:	Signal Transducing Adaptor Molecule 2 (STAM2 Products)
Background:	Hbp, STAM2A, STAM2B, SH3 Domain And ITAM Motif, Hrs-binding protein
Pathways:	EGFR Signaling Pathway, EGFR Downregulation
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL Immunohistochemistry: 5-20 μg/mL Immunocytochemistry: 5-20 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be

handled by trained staff only.

4 °C,-20 °C

24 months


Storage:

Expiry Date:

Storage Comment:


detectable loss of activity. Avoid repeated freeze-thaw cycles.

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without



Immunohistochemistry

Image 1. Detection of STAM2 in Mouse Kidney Tissue using
Polyclonal Antibody to Signal Transducing Adaptor
Molecule 2 (STAM2)



Western Blotting

Image 2. Detection of Recombinant STAM2, Mouse using Polyclonal Antibody to Signal Transducing Adaptor Molecule 2 (STAM2)