



[Go to Product page](#)

Datasheet for ABIN7444523
anti-RIPK2 antibody (AA 432-540)

2 Images

Overview

Quantity:	100 µL
Target:	RIPK2
Binding Specificity:	AA 432-540
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RIPK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Receptor Interacting Serine Threonine Kinase 2 (RIPK2)
Immunogen:	Recombinant Receptor Interacting Serine Threonine Kinase 2 (RIPK2) corresponding to Gln432~Met540 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against RIPK2. It has been selected for its ability to recognize RIPK2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	RIPK2
Alternative Name:	Receptor Interacting Serine Threonine Kinase 2 (RIPK2 Products)
Background:	CCK, CARD3, CARDIAK, GIG30, RICK, RIP2, CARD-containing interleukin-1 beta-converting enzyme-associated kinase, Receptor-interacting 2, RIP-like-interacting CLARP kinase
Pathways:	TCR Signaling , Neurotrophin Signaling Pathway , Activation of Innate immune Response , Cellular Response to Molecule of Bacterial Origin , Positive Regulation of Immune Effector Process , Toll-Like Receptors Cascades

Application Details

Application Notes:	Western blotting: 1-5 µg/mL Immunocytochemistry in formalin fixed cells: 5-20 µg/mL Immunohistochemistry in formalin fixed frozen section: 5-20 µg/mL Immunohistochemistry in paraffin section: 5-20 µg/mL Enzyme-linked Immunosorbent Assay: 0.05-2 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant RIPK2, Human using Polyclonal Antibody to Receptor Interacting Serine Threonine Kinase 2 (RIPK2)