antibodies -online.com









	do to i roddet page

()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL
Target:	EED
Binding Specificity:	AA 142-441
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EED antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Purpose:	Polyclonal Antibody to Embryonic Ectoderm Development (EED)	
Immunogen:	Recombinant Embryonic Ectoderm Development (EED) corresdonding to Asp142~Arg441 with N-terminal His Tag	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against EED. It has been selected for its ability to recognize EED in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Expiry Date:

Target Details	
Target:	EED
Alternative Name:	Embryonic Ectoderm Development (EED Products)
Background:	HEED, WAIT1, WD Protein Associating With Integrin Cytoplasmic Tails 1
Application Details	
Application Notes:	Western blotting: 0.5-2 μ g/mL Immunohistochemistry: 5-20 μ g/mL Immunocytochemistry: 5-20 μ g/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without


detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months



Immunohistochemistry

Image 1. #VALUE!



Western Blotting

Image 2. Detection of Recombinant EED, Human using Polyclonal Antibody to Embryonic Ectoderm Development (EED)