antibodies -online.com





anti-HADH antibody (AA 1-236)



Images



Go to Product pag

()	11	\sim	rv		۱ ۸
	1 \ /	┙	I \/	╙	1/1


Quantity:	100 μL	
Target:	HADH	
Binding Specificity:	AA 1-236	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HADH antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Hydroxyacyl Coenzyme A Dehydrogenase (HADH)		
Immunogen:	Recombinant Hydroxyacyl Coenzyme A Dehydrogenase (HADH) corresdonding to Met1~Arg236		
Isotype:	IgG		
Specificity:	The antibody is a rabbit polyclonal antibody raised against HADH. It has been selected for ability to recognize HADH in immunohistochemical staining and western blotting.		
Cross-Reactivity:	Mouse, Pig, Rat		
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography		


Target Details

. a. got 2 otalio		
Target:	HADH	
Alternative Name:	Hydroxyacyl Coenzyme A Dehydrogenase (HADH Products)	
Background:	HADHSC, HADH1, SCHAD, L-3-Hydroxyacyl-Coenzyme A Dehydrogenase, Short Chain, Medium and short-chain L-3-hydroxyacyl-coenzyme A dehydrogenase	
Pathways:	Negative Regulation of Hormone Secretion, Monocarboxylic Acid Catabolic Process	
Application Details		
Application Notes:	Western blotting: 0.5-2 μ g/mL Immunohistochemistry: 5-20 μ g/mL Immunocytochemistry: 5-20 μ g/mL Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	
	24 months	




Western Blotting

Image 1. Detection of HADH in Porcine Kidney lysate using Polyclonal Antibody to Hydroxyacyl Coenzyme A Dehydrogenase (HADH)



Western Blotting

Image 2. Detection of Recombinant HADH, Human using Polyclonal Antibody to Hydroxyacyl Coenzyme A Dehydrogenase (HADH)



Immunohistochemistry

Image 3. Detection of HADH in Human HepG2 cell using Polyclonal Antibody to Hydroxyacyl Coenzyme A Dehydrogenase (HADH)

Please check the product details page for more images. Overall 6 images are available for ABIN7445138.