
# antibodies -online.com





# anti-EIF4H antibody (AA 2-248)

2 Images



Go to Product page

# Overview

Quantity:	100 μL
Target:	EIF4H
Binding Specificity:	AA 2-248
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF4H antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)


# **Product Details**

Purpose:	Polyclonal Antibody to Eukaryotic Translation Initiation Factor 4H (EIF4H)
Immunogen:	Recombinant Eukaryotic Translation Initiation Factor 4H (EIF4H) corresdonding to Ala2~Glu248 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against EIF4H. It has been selected for its ability to recognize EIF4H in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

# **Target Details**


Target:	EIF4H
Alternative Name:	Eukaryotic Translation Initiation Factor 4H (EIF4H Products)
Background:	WBSCR1, WSCR1, Williams-Beuren Syndrome Chromosome Region 1
Pathways:	SARS-CoV-2 Protein Interactome

Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	1:150-600 Immunohistochemistry: 5-20 µg/mL
	1:15-60 Immunocytochemistry: 5-20 μg/mL
	1:15-60 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



# **Immunohistochemistry**

Image 1. #VALUE!



# **Western Blotting**

**Image 2.** Detection of Recombinant EIF4H, Human using Polyclonal Antibody to Eukaryotic Translation Initiation Factor 4H (EIF4H)