# antibodies -online.com









# Overview

Quantity:	100 μL
Target:	ERK2 (MAPK1)
Binding Specificity:	AA 25-360
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ERK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

# **Product Details**

Purpose:	Polyclonal Antibody to Extracellular Signal Regulated Kinase 2 (ERK2)
Immunogen:	Recombinant Extracellular Signal Regulated Kinase 2 (ERK2) corresdonding to Tyr25~Ser360 with N-terminal His Tag
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ERK2. It has been selected for its ability to recognize ERK2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Chicken, Cow, Dog, Goat, Guinea Pig, Mouse, Pig, Rat, Sheep
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

# **Target Details**

Target:	ERK2 (MAPK1)
Alternative Name:	Extracellular Signal Regulated Kinase 2 (MAPK1 Products)
Background:	MAPK1, P38, P40, ERK2, ERT1, MAPK2, P42MAPK, P42-MAPK, PRKM1, PRKM2, P41, P41mapk, P41-mapk, Mitogen-Activated Protein Kinase 1, Mitogen-activated protein kinase 2
Pathways:	MAPK Signaling, RTK Signaling, Apoptosis, Interferon-gamma Pathway, Fc-epsilon Receptor Signaling Pathway, Response to Growth Hormone Stimulus, Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin, Hepatitis C, Protein targeting to Nucleus, Toll-Like Receptors Cascades, Monocarboxylic Acid Catabolic Process, Autophagy, G-protein mediated Events, Signaling Events mediated by VEGFR1 and VEGFR2, Signaling of Hepatocyte Growth Factor Receptor, VEGFR1 Specific Signals, BCR Signaling, S100 Proteins

# Application Details

Application Notes:	Western blotting: 0.5-2 μg/mL
	1:500-2000 Immunohistochemistry: 5-20 μg/mL
	1:50-200 Immunocytochemistry: 5-20 μg/mL
	1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


# Handling

Format:	Liquid
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Expiry Date:


24 months

# **Images**




### **Immunohistochemistry**

**Image 1.** Detection of ERK2 in Porcine Small intestine Tissue using Polyclonal Antibody to Extracellular Signal Regulated Kinase 2 (ERK2)



# **Western Blotting**

**Image 2.** Detection of Recombinant ERK2, Human using Polyclonal Antibody to Extracellular Signal Regulated Kinase 2 (ERK2)



### **Immunohistochemistry**

Image 3. #VALUE!

Please check the product details page for more images. Overall 7 images are available for ABIN7445753.