

Datasheet for ABIN7446488

Goat anti-Human IgG (Fc Region) Antibody (Biotin)



Overview

| Quantity: | 0.5 mg |
|----------------------|---|
| Target: | IgG |
| Binding Specificity: | Fc Region |
| Reactivity: | Human |
| Host: | Goat |
| Clonality: | Polyclonal |
| Conjugate: | Biotin |
| Application: | ELISA, Immunohistochemistry (IHC), Western Blotting (WB), Immunocytochemistry (ICC) |
| Product Details | |
| Purpose: | F(ab')2 Goat anti-Human IgG-Fc Fragment Antibody Biotinylated |
| Fragment: | F(ab')2 fragment |
| Characteristics: | IgG is the primary immunoglobulin protein found in blood/plasma, and functions to neutralize toxins, active complement, immobilizing pathogens and inducing opsonization, and cell-mediated cytotoxicity. The anti-Fc activity ensures activity only to the Fc portion of the IgG |

Target Details

| Target: | IgG |
|--------------|--------------|
| Abstract: | IgG Products |
| Target Type: | Antibody |

molecule and not the Fab fragments on the light chain.

Application Details

Storage:

Expiry Date:

Storage Comment:

| Application Notes: | IHC: 1:250 - 1:2,500 |
|--------------------|---|
| | ICC: 1:100 - 1:500 |
| | WB: 1:20,000 - 1:400,000 |
| | ELISA: 1:20,000 - 1:400,000 |
| Restrictions: | For Research Use only |
| | |
| Handling | |
| Format: | Liquid |
| Concentration: | 0.5 mg/mL |
| Buffer: | Phosphate Buffered Saline (PBS) containing 0.2 % BSA and 0.09 % Sodium Azide |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which |
| | should be handled by trained staff only. |

4°C

2 - 8°C

12 months