

Datasheet for ABIN744715 anti-PERK antibody (pThr980) (Biotin)



| Overview | |
|----------|--|
| | |

| Quantity: | 100 μL |
|-----------------------|--|
| Target: | PERK (EIF2AK3) |
| Binding Specificity: | pThr980 |
| Reactivity: | Human, Mouse, Rat, Dog, Pig |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PERK antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), |
| | Immunohistochemistry (Frozen Sections) (IHC (fro)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic phosphopeptide derived from mouse PERK around the |
| | phosphorylation site of Thr980 |
| Isotype: | IgG |
| Cross-Reactivity: | Dog, Human, Mouse, Pig, Rat |
| Predicted Reactivity: | Cow,Rabbit |
| Purification: | Purified by Protein A. |
| | |
| Target Details | |
| Target: | PERK (EIF2AK3) |

Target Details

| Alternative Name: | PERK (EIF2AK3 Products) |
|---------------------|--|
| Background: | Synonyms: Pek, Perk, Al427929, Eukaryotic translation initiation factor 2-alpha kinase 3, PRKR- |
| | like endoplasmic reticulum kinase, Pancreatic elF2-alpha kinase, Eif2ak3 |
| | Background: Phosphorylates the alpha subunit of eukaryotic translation-initiation factor 2 |
| | (EIF2), leading to its inactivation and thus to a rapid reduction of translational initiation and |
| | repression of global protein synthesis. Serves as a critical effector of unfolded protein response |
| | (UPR)-induced G1 growth arrest due to the loss of cyclin-D1 (CCND1). Involved in control of |
| | mitochondrial morphology and function. |
| Gene ID: | 13666 |
| UniProt: | Q9Z2B5 |
| Pathways: | Hormone Transport, ER-Nucleus Signaling, Positive Regulation of Endopeptidase Activity, |
| | Hepatitis C, Unfolded Protein Response |
| Application Details | |
| Application Notes: | WB 1:300-5000 |
| | IHC-P 1:200-400 |
| | IHC-F 1:100-500 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Expiry Date: | 12 months |
| | |