# antibodies -online.com





CCR2 Protein-VLP (AA 1-360)





### Overview

Quantity:	100 μg
Target:	CCR2
Protein Characteristics:	AA 1-360
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	VLP
Biological Activity:	Active
Application:	ELISA, Immunogen (Imm), Functional Studies (Func), Surface Plasmon Resonance (SPR)

### **Product Details**

Purpose:	Human CCR2b Protein-VLP
Sequence:	Met1-Leu360
Characteristics:	Recombinant Human CCR2b Protein-VLP is expressed from HEK293.lt contains Met1-Leu360.
Purity:	> 95 % as determined by HPLC
Sterility:	0.22 μm filtered
Endotoxin Level:	Less than 1EU per μg by the LAL method.
Biological Activity Comment:	Immobilized Human CCR2b VLP at 5µg/ml (100µl/Well) on the plate. Dose response curve for
	Anti-CCR2b Antibody, hFc Tag with the EC50 of 12.0ng/ml determined by ELISA.

# **Target Details**

Target:	CCR2
Alternative Name:	CCR2b (CCR2 Products)
Background:	The chemokine (C-C motif) receptor 2B (CCR2B) is one of the two isoforms of the receptor for
	monocyte chemoattractant protein-1 (CCL2), the major chemoattractant for monocytes,
	involved in an array of chronic inflammatory diseases. The actin-binding protein filamin A
	(FLNa) as a protein that associates with the carboxyl-terminal tail of CCR2B. FLNa emerges as
	an important protein for controlling the internalization and spatial localization of the CCR2B
	receptor in different dynamic membrane structures.
Molecular Weight:	42.9kDa.
Pathways:	cAMP Metabolic Process, Regulation of Leukocyte Mediated Immunity, Positive Regulation of
	Immune Effector Process
Application Details	
Application Notes:	Antibody Discovery: Immunization, Screening, Functional Characterization
	Affinity determination: ELISA, SPR
	In vivo pharmacokinetic analysis
	CMC method development
	CAR-T Positive Rate Detection Plead cample determination: ELISA
	Blood sample determination: ELISA
Comment:	Virus-like particles (VLPs) are formed from the outer capsid protein of a virus and are tiny
	nanoparticles formed by the automatic assembly of one or more capsid proteins. VLPs do not
	contain viral infectious genomes, so they are relatively safe during production operations. The
	SAMS <sup>™</sup> protein engineering platform has been used to express a series of biotinylated, non-
	biotinylated, and fluorescently-labeled VLP-displayed antigens. They are suitable for SPR, ELISA
	CAR-T positive rate detection, and other experimental scenarios.
	Virus-Like Particles (VLPs) are highly immunogenic, meaning that they can elicit a strong
	immune response in the host. VLPs are recognized by the immune system and are taken up by
	antigen-presenting cells (APCs) such as dendritic cells. Once taken up by APCs, VLPs are
	processed and presented to T cells, which can trigger the activation of B cells to produce
	antibodies against the displayed antigen. Because VLPs resemble the structure and
	composition of native viruses, they are highly effective at inducing both humoral and cellular .
	immune responses.

# **Application Details**

Generally, VLPs range in size from approximately 20 to 200 nanometers (nm). Compared to a
cell-based immunization approach, their smaller size can optimize the immune response to
target the specific antigen displayed on the surface of the engineered VLPs.


Restrictions:

For Research Use only

# Handling

Format:	Liquid
Buffer:	Supplied as 0.22µm filtered solution in PBS (pH 7.4). Notice: If you need it for immunization, Do Not use any adjuvant.
Storage:	-80 °C
Storage Comment:	Valid for 12 months from date of receipt when stored at -80°C., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Expiry Date:	12 months

## **Images**




# Size-exclusion chromatography-High Pressure Liquid Chromatography

**Image 1.** The purity of Human CCR2b VLP is greater than 95 % as determined by SEC-HPLC.

#### **Human CCR2b VLP ELISA**

0.5µg Human CCR2b VLP Per Well



Log Anti-CCR2b Antibody, hFc Tag Conc.(µg/ml)

### **ELISA**

Image 2. Immobilized Human CCR2b VLP at 5  $\mu$ g/mL (100  $\mu$ L/Well) on the plate. Dose response curve for Anti-CCR2b Antibody, hFc Tag with the EC50 of 12.0 ng/mL determined by ELISA.