

Datasheet for ABIN7448169

GPRC5D Protein-VLP (AA 1-345)[Go to Product page](#)**2** Images

Overview

| | |
|--------------------------|--|
| Quantity: | 100 µg |
| Target: | GPRC5D |
| Protein Characteristics: | AA 1-345 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | VLP |
| Biological Activity: | Active |
| Application: | ELISA, Functional Studies (Func), Immunogen (Imm), Surface Plasmon Resonance (SPR) |

Product Details

| | |
|------------------------------|---|
| Purpose: | Human GPRC5D Protein-VLP |
| Sequence: | Met1-Val345 |
| Characteristics: | Recombinant Human GPRC5D Protein-VLP is expressed from HEK293. It contains Met1-Val345. |
| Purity: | > 95 % as determined by HPLC |
| Sterility: | 0.22 µm filtered |
| Endotoxin Level: | Less than 1EU per µg by the LAL method. |
| Biological Activity Comment: | Immobilized Human GPRC5D VLP at 5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-GPRC5D Antibody, hFc Tag with the EC50 of 11.8ng/ml determined by ELISA. |

Target Details

| | |
|-------------------|--|
| Target: | GPRC5D |
| Alternative Name: | GPRC5D (GPRC5D Products) |
| Background: | Chimeric antigen receptor (CAR) T cells, a type of cell-based immunotherapy, have shown some promising results in multiple myeloma, a bone marrow cancer. The orphan G protein-coupled receptor, class C group 5 member D (GPRC5D), normally expressed only in the hair follicle, Using quantitative immunofluorescence, we determined that GPRC5D protein is expressed on CD138 MM cells from primary marrow samples with a distribution that was similar to, but independent of, BCMA. |
| Molecular Weight: | 39.6 kDa. |


Application Details

| | |
|--------------------|---|
| Application Notes: | <ul style="list-style-type: none">• Antibody Discovery: Immunization, Screening, Functional Characterization• Affinity determination: ELISA, SPR• In vivo pharmacokinetic analysis• CMC method development• CAR-T Positive Rate Detection• Blood sample determination: ELISA |
| Comment: | <p>Virus-like particles (VLPs) are formed from the outer capsid protein of a virus and are tiny nanoparticles formed by the automatic assembly of one or more capsid proteins. VLPs do not contain viral infectious genomes, so they are relatively safe during production operations. The SAMS™ protein engineering platform has been used to express a series of biotinylated, non-biotinylated, and fluorescently-labeled VLP-displayed antigens. They are suitable for SPR, ELISA, CAR-T positive rate detection, and other experimental scenarios.</p> <p>Virus-Like Particles (VLPs) are highly immunogenic, meaning that they can elicit a strong immune response in the host. VLPs are recognized by the immune system and are taken up by antigen-presenting cells (APCs) such as dendritic cells. Once taken up by APCs, VLPs are processed and presented to T cells, which can trigger the activation of B cells to produce antibodies against the displayed antigen. Because VLPs resemble the structure and composition of native viruses, they are highly effective at inducing both humoral and cellular immune responses.</p> <p>Generally, VLPs range in size from approximately 20 to 200 nanometers (nm). Compared to a cell-based immunization approach, their smaller size can optimize the immune response to</p> |

Application Details


| | |
|------------------|---|
| | target the specific antigen displayed on the surface of the engineered VLPs. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | Supplied as 0.22µm filtered solution in PBS (pH 7.4). Notice: If you need it for immunization, Do Not use any adjuvant. |
| Storage: | -80 °C |
| Storage Comment: | Valid for 12 months from date of receipt when stored at -80°C., Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Expiry Date: | 12 months |

Images



Size-exclusion chromatography-High Pressure Liquid Chromatography

Image 1. The purity of Human GPRC5D VLP is greater than 95 % as determined by SEC-HPLC.



ELISA

Image 2. Immobilized Human GPRC5D VLP at 5 µg/mL (100 µL/Well) on the plate. Dose response curve for Anti-GPRC5D Antibody, hFc Tag with the EC50 of 11.8 ng/mL determined by ELISA.