

Datasheet for ABIN744840

anti-Phospholipase C gamma 2 antibody (pTyr1217) (FITC)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Phospholipase C gamma 2 (PLCG2)
Binding Specificity:	pTyr1217
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Phospholipase C gamma 2 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human PLC gamma 2 around the phosphorylation site of Tyr1217
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	Phospholipase C gamma 2 (PLCG2)
---------	---------------------------------

Target Details

Alternative Name:	PLCG2 (PLCG2 Products)
Background:	Synonyms: PLCG2phospho Y1217, PLC _2Tyr1217, PLC gamma 2/PLCG2Tyr1217, PLC 2, PLC gamma 2, PLC IV, PLCG2, 1 phosphatidylinositol 4 5 biphosphate phosphodiesterase gamma 2, EC 3.1.4.11, Phosphoinositide phospholipase C, Phospholipase C gamma 2, Phospholipase C, gamma 2 phosphatidylinositol specic, PLCG2_HUMAN. Background: Enzymes of the phospholipase C family catalyze the hydrolysis of phospholipids to yield diacylglycerols and water soluble phosphorylated derivatives of the lipid head groups. A number of these enzymes have specificity for phosphoinositides. Of the phosphoinositide specific phospholipase C enzymes, C beta is regulated by heterotrimeric G protein coupled receptors, while the closely related C gamma 1 and C gamma 2 enzymes are controlled by receptor tyrosine kinases. The C gamma 1 and C gamma 2 enzymes are composed of phospholipase domains that flank regions of homology to noncatalytic domains of the SRC oncogene product, SH2 and SH3.
Gene ID:	5336
Pathways:	RTK Signaling , WNT Signaling , Fc-epsilon Receptor Signaling Pathway , Inositol Metabolic Process , Myometrial Relaxation and Contraction , Toll-Like Receptors Cascades , VEGF Signaling , BCR Signaling

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months