

## Datasheet for ABIN7450611 anti-PSMD2 antibody (AA 1-50)



## Go to Product page

$\sim$				
$O_1$	<b>/</b> el	rVI	(0)	Λ

Background:

Quantity:	100 μg
Target:	PSMD2
Binding Specificity:	AA 1-50
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMD2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP)
Product Details	
Purpose:	Rabbit anti-PSMD2 Antibody, Affinity Purified
Immunogen:	Between AA 1 and 50
Isotype:	IgG
Predicted Reactivity:	Orangutan
Purification:	Affinity Purified
Target Details	
Target:	PSMD2
Alternative Name:	PSMD2 (PSMD2 Products)

Background: PSMD2 [proteasome (prosome, macropain) 26S subunit, non-ATPase, 2] is a

subunit of the 26 S proteasome. The 26S proteasome is a multicatalytic proteinase complex
with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The
20S core is composed of 4 rings of 28 non-identical subunits, 2 rings are composed of 7 alpha
subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a
base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains
up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a
high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-
lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome,
is the processing of class I MHC peptides. PSMD2 is a subunit of the 19S regulator lid, which is
responsible for substrate recognition and binding [taken from NCBI Entrez Gene (Gene ID:
5708)].

	Gene ID:	5708	
--	----------	------	--

NP\_002799

UniProt:	Q13200

Pathwavs:	Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA, Ubiquitin Proteasome Pathway
r attivayo.	Time to or or nacco, but the phoation, cynthesis or but, conquiting receasing rathway

## **Application Details**

NCBI Accession:

Application Notes:	IP: 2 - 10 μg/mg lysate
	WB: 1:2,000 - 1:10,000

Restrictions: For Research Use only

## Handling

Concentration:	1000 μg/mL
Buffer:	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months