

Datasheet for ABIN7451264
anti-IDH1 antibody (AA 1-50)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	IDH1
Binding Specificity:	AA 1-50
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IDH1 antibody is un-conjugated
Application:	Immunoprecipitation (IP)

Product Details

Purpose:	Rabbit anti-IDH1 Antibody, Affinity Purified
Immunogen:	Between AA 1 and 50
Isotype:	IgG
Predicted Reactivity:	Bovine,Orangutan
Purification:	Affinity Purified

Target Details

Target:	IDH1
Alternative Name:	IDH1 (IDH1 Products)
Background:	Background: Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to

Target Details

2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. IDH1 is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production [taken from NCBI Entrez Gene (Gene ID: 3417)].

Gene ID:	3417
NCBI Accession:	NP_005887
UniProt:	O75874
Pathways:	Warburg Effect

Application Details

Application Notes:	IP: 2 - 10 µg/mg lysate WB: Not recommended. Use rabbit anti-IDH1 antibody ABIN7450997.
Restrictions:	For Research Use only

Handling

Concentration:	1000 µg/mL
Buffer:	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months