antibodies -online.com





anti-MED15 antibody (AA 650-700)



\sim	
()\/\	rview
\cup	1 410 44

Quantity:	20 μg
Target:	MED15
Binding Specificity:	AA 650-700
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED15 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP)

Product Details

Purpose:	Rabbit anti-MED15 Antibody, Affinity Purified
Immunogen:	between AA 650 and 700
Isotype:	IgG
Predicted Reactivity:	Mouse
Purification:	Affinity Purified

Target Details

Target:	MED15
Alternative Name:	MED15 (MED15 Products)
Background:	Background: Mediator complex subunit 15 is a component of the Mediator complex, a

coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. MED15 is required for cholesterol-dependent gene regulation. It also positively regulates the Nodal signaling pathway. [taken from the Universal Protein Resource (UniProt) Q96RN5].

Gene ID: 51586

NCBI Accession: NP_001003891

UniProt: Q96RN5

Pathways: Stem Cell Maintenance, Regulation of Lipid Metabolism by PPARalpha

Application Details

Application Notes: IP: $2 - 5 \mu g/mg$ lysate

WB: 1:2,000 - 1:10,000

Restrictions: For Research Use only

Handling

Concentration:	200 μg/mL
Buffer:	Tris-buffered Saline containing 0.1 % BSA and 0.09 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Expiry Date:	12 months